

MEETING ABSTRACT

Open Access

# Viral DNA replication: new insights and discoveries from large scale computational analysis

Darius Kazlauskas\*, Česlovas Venclovas

From Third International Society for Computational Biology (ISCB) European Student Council Symposium 2014  
Strasbourg, France. 6 September 2014

## Background

The ability to replicate is essential for all living entities. Duplication of genetic information is carried out by replication proteins. DNA replication has been well studied in T7, T4 phages and herpes viruses; however, the information about replication mechanisms from other groups of viruses is either scarce or missing altogether. Double-stranded (ds) DNA viruses infect cells from all domains of life, they evolve fast and are very diverse. Their genome size varies from 5 to 2,500 kbp.

Published: 13 February 2015

doi:10.1186/1471-2105-16-S3-A2

Cite this article as: Kazlauskas and Venclovas: Viral DNA replication: new insights and discoveries from large scale computational analysis. *BMC Bioinformatics* 2015 **16**(Suppl 3):A2.

## Results and conclusions

To better understand viral DNA replication, we identified replication proteins in dsDNA viruses using current state-of-the-art homology detection methods. Over 150,000 proteins from 1,574 genomes were analyzed. We found that the composition of replication machinery depends on the virus genome size. Small viruses (<40 kbp) use protein-primed DNA replication or rely on replication proteins from the host. Large viruses (>140 kbp) have their own RNA-primed replication apparatus often supplemented with processivity factors and DNA topoisomerases to increase replication speed and efficiency. This insight led us to a search for „missing“ replication components in large genomes and resulted in the discovery of single-stranded DNA binding (SSB) proteins in larger eukaryotic viruses. Surprisingly these proteins turned out to be homologs of SSB proteins previously thought to be specific for T7-like phages. Additionally with the analysis of the herpes viral helicase-primase complex we found that one of its components, UL8, is a highly diverged inactivated B-family DNA polymerase.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



Institute of Biotechnology, Vilnius University, Lithuania